*Neuroscience Letters*, 142 (1992) 275–278
Elsevier Scientific Publishers Ireland Ltd.



## AUTHOR INDEX

Aitkin, L. and Jones, R., Azimuthal processing in the posterior auditory thalamus of cats, 81
Akiguchi, I., see Nagao, M., 131
Akutsu, S., Takada, M., Ohki-Hamazaki, H., Murakami, S. and Arai, Y., Origin of luteinizing hormone-releasing hormone (LHRH) neurons in the chick embryo: effect of the olfactory placode ablation, 241
Alcantara, S., see Soriano, E., 147
Allsop, D., see Gentleman, S.M., 9
Alonso-Bedate, M., see Alonso-Gómez, A.L., 187
Alonso-Gómez, A.L., Alonso-Bedate, M. and Delgado, M.J., Thermal sensitivity and effect of temperature acclimation on ocular serotonin *N*-acetyltransferase activity in *Rana perezi*, 187
Åman, K., see Hökfelt, T., 139
Aoki, T., see Kamei, J., 13
Arai, Y., see Akutsu, S., 241
Arvidsson, U., see Hökfelt, T., 139
Auladell, C., see Soriano, E., 147

Bakay, R.A.E., see Kennedy, P.R., 89
Baldauf, J., see Jurna, I., 62
Baram, T.Z. and Schultz, L., CRH gene expression in the fetal rat is not increased after pharmacological adrenalectomy, 215
Bartfai, T., see Hökfelt, T., 139
Batchelor, A., see Hansel, C., 211
Beach, T.G. and McGeer, E.G., Cholinergic fiber loss occurs in the absence of synaptophysin depletion in Alzheimer's disease primary visual cortex, 253
Beart, P.M., Mercer, L.D., Conway, E.L. and Searle, A.J., Polyamine-sensitive binding of [$^{125}$I]ifenprodil in washed, frozen-thawed synaptic membranes: evidence for high affinity binding requiring an open NMDA channel, 219
Bedecs, K., see Hökfelt, T., 139
Berger, M.L., Seifriz, I., Letschnig, M., Schödl, C. and Noe, C.R., Interaction of long chain n-alkyl diamines with the NMDA receptor complex, 85
Bernardi, G., see Calabresi, P., 95
Bernau, N.A., see Pubols, L.M., 111
Biggs, C.S., see Whitton, P.S., 5
Bogónez, E., Gómez-Puertas, P. and Satrústegui, J., Pyruvate dehydrogenase dephosphorylation in rat brain synaptosomes and mitochondria: evidence for a calcium-mediated effect in response to depolarization, and variations due to ageing, 123
Bravo, R., see Herdegen, T., 57
Brouwer, N., Van Dijken, H., Ruiters, M.H.J., Van Willigen, J.-D. and Horst, G.J.T., Localization of dopamine $D_2$ receptor mRNA with non-radioactive in situ hybridization histochemistry, 223
Buckmaster, P.S., see Strowbridge, B.W., 205
Bunnemann, B., Iwai, N., Metzger, R., Fuxe, K., Inagami, T. and Ganten, D., The distribution of angiotensin II $AT_1$ receptor subtype mRNA in the rat brain, 155
Bureš, J., see Engelmann, M., 69
Burgoyne, R.D., see Pearson, H., 27
Burleigh, A.L., see Pubols, L.M., 111

Calabresi, P., Maj, R., Mercuri, N.B. and Bernardi, G., Coactivation of $D_1$ and $D_2$ dopamine receptors is required for long-term synaptic depression in the striatum, 95
Cameron, D.L., see Crocker, A.D., 73
Ceccatelli, S., see Hökfelt, T., 139
Cheng, K.W., A heparin-binding protein in porcine seminal plasma stimulates neurite outgrowth on neuroblastoma cells in culture, 77
Chrapusta, S. and Grieb, P., Evidence for sex steroid receptors in feline brainstem, 167
Clifford, D.B., see Izumi, Y., 163
Clinton, J., see Gentleman, S.M., 9
Close, S.P., see Elliott, P.J., 1
Conway, E.L., see Beart, P.M., 219
Costentin, J., see Kitabgi, P., 200
Cremel, G., see Laabich, A., 196
Crocker, A.D. and Cameron, D.L., Localisation of striatal muscarinic receptors involved in dopamine receptor-mediated behavioural responses, 73
Cuber, J.-C., see Kitabgi, P., 200
Cuénod, M., see Hansel, C., 211

Dawson, D.A., Kusumoto, K., Graham, D.I., McCulloch, J. and Macrae, I.M., Inhibition of nitric oxide synthesis does not reduce infarct volume in a rat model of focal cerebral ischaemia, 151
Dawson, S.D., see Pubols, L.M., 111
De Lecea, L., see Soriano, E., 147
Del Rio, J.A., see Soriano, E., 147
Delaunoy, J.P., see Laabich, A., 196
Delgado, M.J., see Alonso-Gómez, A.L., 187
Do, K.Q., see Hansel, C., 211
Doulut, S., see Kitabgi, P., 200
Dubuc, I., see Kitabgi, P., 200

Elde, R., see Wu, W., 143
Elliott, P.J., Walsh, D.M. and Close, S.P., Dopamine $D_1$ and $D_2$ receptor interactions in the MPTP-treated marmoset, 1
Endo, K., see Hori, Y., 191
Engelmann, M., Bureš, J. and Landgraf, R., Vasopressin administration via microdialysis into the septum interferes with the acquisition of spatial memory in rats, 69

Ferrer, I., see Soriano, E., 147
Fiallos-Estrada, C., see Herdegen, T., 57
Flaherty, K.N., see Greif, K.F., 22
Fowler, L.J., see Whitton, P.S., 5
Fujimoto, K. and Kita, H., Responses of rat substantia nigra pars reticulata units to cortical stimulation, 105
Fulcrand, H., see Kitabgi, P., 200
Fuxe, K., see Bunnemann, B., 155

Ganten, D., see Bunnemann, B., 155
García-Sevilla, J.A., see Pascual, J., 36
Garthwaite, J., see Hansel, C., 211
Gentleman, S.M., Williams, B., Royston, M.C., Jagoe, R., Clinton, J., Perry, R.H., Ince, P.G., Allsop, D., Polak, J.M. and Roberts, G.W., Quantification of $\beta$A4 protein deposition in the medial temporal

lobe: a comparison of Alzheimer's disease and senile dementia of the Lewy body type, 9
Glenfield, P.J., see Heath, J.W., 17
Gómez-Puertas, P., see Bogónez, E., 123
Graham, D.I., see Dawson, D.A., 151
Graham, M.E., see Pearson, H., 27
Greif, K.F. and Flaherty, K.N., Age-dependent effects of deafferentation of the rat superior cervical ganglion on expression of P65 (synaptotagmin) during postnatal development, 22
Grieb, P., see Chrapusta, S., 167
Grijalba, B., see Pascual, J., 36

Hansel, C., Batchelor, A., Cuénod, M., Garthwaite, J., Knöpfel, T. and Do, K.Q., Delayed increase of extracellular arginine, the nitric oxide precursor, following electrical white matter stimulation in rat cerebellar slices, 211
Heath, J.W., Glenfield, P.J. and Rostas, J.A.P., Structural maturation of synapses in the rat superior cervical ganglion continues beyond four weeks of age, 17
Herdegen, T., Fiallos-Estrada, C., Schmid, W., Bravo, R. and Zimmermann, M., The transcription factor CREB, but not immediate-early gene encoded proteins, is expressed in activated microglia of lumbar spinal cord following sciatic nerve transection in the rat, 57
Hökfelt, T., Åman, K., Arvidsson, U., Bedecs, K., Ceccatelli, S., Hulting, A.-L., Langel, U., Meister, B., Pieribone, V. and Bartfai, T., Galanin message-associated peptide (GMAP)- and galanin-like immunoreactivities: overlapping and differential distributions in the rat, 139
Hökfelt, T., see Wu, W., 143
Hori, Y. and Endo, K., Miniature postsynaptic currents recorded from identified rat spinal dorsal horn projection neurons in thin-slice preparations, 191
Horst, G.J.T., see Brouwer, N., 223
Houtani, T., see Sato, K., 48
Hulting, A.-L., see Hökfelt, T., 139

Ikeda, J., Mies, G., Nowak Jr., T.S., Joó, F. and Klatzo, I., Evidence for increased calcium influx across the choroid plexus following brief ischemia of gerbil brain, 257
Ikeda, M., see Sato, K., 48
Imamura, K., see Mataga, N., 115
Inagami, T., see Bunnemann, B., 155
Ince, P.G., see Gentleman, S.M., 9
Inoki, R., see Yonehara, N., 53
Iwai, N., see Bunnemann, B., 155
Izumi, H., see Kuchiiwa, S., 237
Izumi, Y., Clifford, D.B. and Zorumski, C.F., Norepinephrine reverses *N*-methyl-D-aspartate-mediated inhibition of long-term potentiation in rat hippocampal slices, 163

Jagoe, R., see Gentleman, S.M., 9
Jellinger, K., see Sofic, E., 128
Jones, R., see Aitkin, L., 81
Joó, F., see Ikeda, J., 257
Jurna, I., Kaiser, R., Kretz, O. and Baldauf, J., Oral naloxone reduces constipation but not antinociception from oral morphine in the rat, 62

Kaiser, R., see Jurna, I., 62
Kamei, J., Aoki, T. and Kasuya, Y., Periaqueductal gray matter stimulation-produced analgesia in diabetic rats, 13
Kamo, H., see Nagao, M., 131
Kane, L.A., see Pubols, L.M., 111
Karita, K., see Kuchiiwa, S., 237
Kasuya, Y., see Kamei, J., 13
Katsuki, H., Saito, H. and Satoh, M., The involvement of muscarinic, $\beta$-adrenergic and metabotropic glutamate receptors in long-term potentiation in the fimbria–CA3 pathway of the hippocampus, 249
Kawata, M., see Yuri, K., 135
Kennedy, P.R., Mirra, S.S. and Bakay, R.A.E., The cone electrode: ultrastructural studies following long-term recording in rat and monkey cortex, 89
Kimura, J., see Nagao, M., 131
Kita, H., see Fujimoto, K., 105
Kitabgi, P., Dubuc, I., Nouel, D., Costentin, J., Cuber, J.-C., Fulcrand, H., Doulut, S., Rodriguez, M. and Martinez, J., Effects of thiorphan, bestatin and a novel metallopeptidase inhibitor JMV 390-1 on the recovery of neurotensin and neuromedin N released from mouse hypothalamus, 200
Klatzo, I., see Ikeda, J., 257
Knöpfel, T., see Hansel, C., 211
Kogure, K., see Yamasaki, Y., 45
Kondo, H., see Toyoda, J., 41
Konishi, S., Song, S.-Y. and Saito, K., An after-depolarization following action potentials and its modulation by substance P in rat sympathetic neurons, 245
Kretz, O., see Jurna, I., 62
Kuchiiwa, S., Izumi, H., Karita, K. and Nakagawa, S., Origins of parasympathetic postganglionic vasodilator fibers supplying the lips and gingivae; an WGA–HRP study in the cat, 237
Kujiraoka, T., see Toyoda, J., 41
Kumazawa, T., see Sato, K., 48
Kunow, M., see Wachtel, H., 179
Kusumoto, K., see Dawson, D.A., 151

Laabich, A., Delaunoy, J.P., Cremel, G. and Staedel, C., Modifications of ependymal cells membranes by galactocerebrosides in cell culture, 196
Landgraf, R., see Engelmann, M., 69
Lange, K.W., see Sofic, E., 128
Langel, U., see Hökfelt, T., 139
Lartius, R., see Uemura, E., 171
LeDoux, J.E., see Romanski, L.M., 228
Leigh, R.J., see Ott, D., 183
Letschnig, M., see Berger, M.L., 85
Liu, R.-H. and Zhao, Z.-Q., Selective blockade by yohimbine of descending spinal inhibition from lateral reticular nucleus but not from locus coeruleus in rats, 65
Löschmann, P.-A., see Wachtel, H., 179
Lynch, G.R., see Margraf, R.R., 175

Macrae, I.M., see Dawson, D.A., 151
Mailleux, P., Vanderwinden, J.-M. and Vanderhaeghen, J.-J., The new growth factor pleiotrophin (HB-GAM) mRNA is selectively present in the meningothelial cells of human meningiomas, 31
Maj, R., see Calabresi, P., 95
Margraf, R.R., Puchalski, W. and Lynch, G.R., Absence of a daily neuronal rhythm in the suprachiasmatic nuclei of acircadian Djungarian hamsters, 175
Martinez, J., see Kitabgi, P., 200
Mataga, N., Imamura, K. and Watanabe, Y., L-*threo*-3,4-Dihydroxyphenylserine enhanced ocular dominance plasticity in adult cats, 115
Matsuura, H., see Yonehara, N., 53

Matsuura, N., see Yamasaki, Y., 45
McCulloch, J., see Dawson, D.A., 151
McGeer, E.G., see Beach, T.G., 253
Meister, B., see Hökfelt, T., 139
Mercer, L.D., see Beart, P.M., 219
Mercuri, N.B., see Calabresi, P., 95
Metzger, R., see Bunnemann, B., 155
Mies, G., see Ikeda, J., 257
Minachi, M., see Uemura, E., 171
Mirra, S.S., see Kennedy, P.R., 89
Mori, M., see Okada, Y., 233
Murakami, S., see Akutsu, S., 241
Murray, C.E., see See, R.E., 100

Nagao, M., Kamo, H., Akiguchi, I. and Kimura, J., Soybean agglutinin binds commonly to a subpopulation of small-diameter neurons in dorsal root ganglion, vascular endothelium and microglia in human spinal cord, 131
Nakagawa, S., see Kuchiiwa, S., 237
Noe, C.R., see Berger, M.L., 85
Nouel, D., see Kitabgi, P., 200
Nowak Jr., T.S., see Ikeda, J., 257

Ohki-Hamazaki, H., see Akutsu, S., 241
Oka, Y., Gonadotropin-releasing hormone (GnRH) cells of the terminal nerve as a model neuromodulator system, 119
Okada, Y., Sakurai, T. and Mori, M., Excitatory effect of adenosine on neurotransmission is due to increase of transmitter release in the hippocampal slices, 233
Onodera, H., see Yamasaki, Y., 45
Ott, D., Seidman, S.H. and Leigh, R.J., The stability of human eye orientation during visual fixation, 183

Pascual, J., Grijalba, B., García-Sevilla, J.A., Zarranz, J.J. and Pazos, A., Loss of high-affinity $\alpha_2$-adrenoceptors in Alzheimer's disease: an autoradiographic study in frontal cortex and hippocampus, 36
Pazos, A., see Pascual, J., 36
Pearce, B.R., see Whitton, P.S., 5
Pearson, H., Graham, M.E. and Burgoyne, R.D., *N*-Methyl-D-aspartate responses in rat cerebellar granule cells are modified by chronic depolarisation in culture, 27
Perry, R.H., see Gentleman, S.M., 9
Pieribone, V., see Hökfelt, T., 139
Polak, J.M., see Gentleman, S.M., 9
Polans, A.S., see Pubols, L.M., 111
Pubols, L.M., Bernau, N.A., Kane, L.A., Dawson, S.D., Burleigh, A.L. and Polans, A.S., Distribution of 5-$HT_1$ binding sites in cat spinal cord, 111
Puchalski, W., see Margraf, R.R., 175

Riederer, P., see Sofic, E., 128
Roberts, G.W., see Gentleman, S.M., 9
Rodriguez, M., see Kitabgi, P., 200
Romanski, L.M. and LeDoux, J.E., Bilateral destruction of perirhinal and neocortical projection targets of the acoustic thalamus does not disrupt auditory fear conditioning, 228
Rostas, J.A.P., see Heath, J.W., 17
Royston, M.C., see Gentleman, S.M., 9
Ruiters, M.H.J., see Brouwer, N., 223

Saito, H., see Katsuki, H., 249
Saito, K., see Konishi, S., 245
Sakurai, T., see Okada, Y., 233
Sato, K., Houtani, T., Ueyama, T., Ikeda, M., Yamashita, T., Kumazawa, T. and Sugimoto, T., Mapping of the cochlear nucleus subregions in the rat with neuronal Fos protein induced by acoustic stimulation with low tones, 48
Satoh, M., see Katsuki, H., 249
Satrústegui, J., see Bogónez, E., 123
Sawada, T., see Yonehara, N., 53
Schmid, W., see Herdegen, T., 57
Schödl, C., see Berger, M.L., 85
Schultz, L., see Baram, T.Z., 215
Schwartzkroin, P.A., see Strowbridge, B.W., 205
Searle, A.J., see Beart, P.M., 219
Seckl, J.R., see Yau, J.L.W., 159
See, R.E. and Murray, C.E., Changes in striatal dopamine release and metabolism during and after subchronic haloperidol administration in rats, 100
Seidman, S.H., see Ott, D., 183
Seifriz, I., see Berger, M.L., 85
Shimbo, K., see Toyoda, J., 41
Sofic, E., Lange, K.W., Jellinger, K. and Riederer, P., Reduced and oxidized glutathione in the substantia nigra of patients with Parkinson's disease, 128
Song, S.-Y., see Konishi, S., 245
Soriano, E., Del Rio, J.A., Ferrer, I., Auladell, C., De Lecea, L. and Alcantara, S., Late appearance of parvalbumin-immunoreactive neurons in the rodent cerebral cortex does not follow an 'inside-out' sequence, 147
Staedel, C., see Laabich, A., 196
Strowbridge, B.W., Buckmaster, P.S. and Schwartzkroin, P.A., Potentiation of spontaneous synaptic activity in rat mossy cells, 205
Sugimoto, T., see Sato, K., 48
Suzuki, T., see Yamasaki, Y., 45

Takada, M., see Akutsu, S., 241
Toyoda, J., Shimbo, K., Kondo, H. and Kujiraoka, T., Push-pull modulation of ganglion cell responses of carp retina by amacrine cells, 41

Uemura, E., Minachi, M. and Lartius, R., Enhanced neurite growth in cultured neuroblastoma cells exposed to aluminum, 171
Ueyama, T., see Sato, K., 48

Van Dijken, H., see Brouwer, N., 223
Van Willigen, J.-D., see Brouwer, N., 223
Vanderhaeghen, J.-J., see Mailleux, P., 31
Vanderwinden, J.-M., see Mailleux, P., 31

Wachtel, H., Kunow, M. and Löschmann, P.-A., NBQX (6-nitro-sulfamoyl-benzo-quinoxaline-dione) and CPP (3-carboxy-piperazin-propyl phosphonic acid) potentiate dopamine agonist induced rotations in substantia nigra lesioned rats, 179
Walsh, D.M., see Elliott, P.J., 1
Watanabe, Y., see Mataga, N., 115
Wessendorf, M.W., see Wu, W., 143
Whitton, P.S., Biggs, C.S., Pearce, B.R. and Fowler, L.J., Regional effects of MK-801 on dopamine and its metabolites studied by in vivo microdialysis, 5
Williams, B., see Gentleman, S.M., 9
Wu, W., Elde, R., Wessendorf, M.W. and Hökfelt, T., Identification of neurons expressing thyrotropin releasing-hormone receptor mRNA in spinal cord and lower brainstem of rat, 143

Yamasaki, Y., Suzuki, T., Yamaya, H., Matsuura, N., Onodera, H. and Kogure, K., Possible involvement of interleukin-1 in ischemic brain edema formation, 45
Yamashita, T., see Sato, K., 48
Yamaya, H., see Yamasaki, Y., 45
Yau, J.L.W. and Seckl, J.R., Central 6-hydroxydopamine lesions decrease mineralocorticoid, but not glucocorticoid receptor gene expression in the rat hippocampus, 159
Yonehara, N., Sawada, T., Matsuura, H. and Inoki, R., Influence of electro-acupuncture on the release of substance P and the potential evoked by tooth pulp stimulation in the trigeminal nucleus caudalis of the rabbit, 53
Yuri, K. and Kawata, M., Nuclear localization of estrogen receptor-immunoreactivity in the preoptic area of female rats and its reduction by intraventricular colchicine treatment, 135

Zarranz, J.J., see Pascual, J., 36
Zhao, Z.-Q., see Liu, R.-H., 65
Zimmermann, M., see Herdegen, T., 57
Zorumski, C.F., see Izumi, Y., 163

---

**CONTENTS** *(continued from the back cover)*

Effects of thiorphan, bestatin and a novel metallopeptidase inhibitor JMV 390-1 on the recovery of neurotensin and neuromedin N released from mouse hypothalamus
P. Kitabgi, I. Dubuc, D. Nouel, J. Costentin, J.-C. Cuber, H. Fulcrand, S. Doulut, M. Rodriguez and J. Martinez (France) 200
Potentiation of spontaneous synaptic activity in rat mossy cells
B.W. Strowbridge, P.S. Buckmaster and P.A. Schwartzkroin (USA) 205
Delayed increase of extracellular arginine, the nitric oxide precursor, following electrical white matter stimulation in rat cerebellar slices
C. Hansel, A. Batchelor, M. Cuénod, J. Garthwaite, T. Knöpfel and K.Q. Do (Switzerland, UK) 211
CRH gene expression in the fetal rat is not increased after pharmacological adrenalectomy
T.Z. Baram and L. Schultz (USA) 215
Polyamine-sensitive binding of [$^{125}$I]ifenprodil in washed, frozen-thawed synaptic membranes: evidence for high affinity binding requiring an open NMDA channel
P.M. Beart, L.D. Mercer, E.L. Conway and A.J. Searle (Australia) 219
Localization of dopamine $D_2$ receptor mRNA with non-radioactive in situ hybridization histochemistry
N. Brouwer, H. Van Dijken, M.H.J. Ruiters, J.-D. Van Willigen and G.J.T. Horst (Netherlands) 223
Bilateral destruction of perirhinal and neocortical projection targets of the acoustic thalamus does not disrupt auditory fear conditioning
L.M. Romanski and J.E. LeDoux (USA) 228
Excitatory effect of adenosine on neurotransmission is due to increase of transmitter release in the hippocampal slices
Y. Okada, T. Sakurai and M. Mori (Japan) 233
Origins of parasympathetic postganglionic vasodilator fibers supplying the lips and gingivae; an WGA–HRP study in the cat
S. Kuchiiwa, H. Izumi, K. Karita and S. Nakagawa (Japan) 237
Origin of luteinizing hormone-releasing hormone (LHRH) neurons in the chick embryo: effect of the olfactory placode ablation
S. Akutsu, M. Takada, H. Ohki-Hamazaki, S. Murakami and Y. Arai (Japan) 241
An after-depolarization following action potentials and its modulation by substance P in rat sympathetic neurons
S. Konishi, S.-Y. Song and K. Saito (Japan) 245
The involvement of muscarinic, $\beta$-adrenergic and metabotropic glutamate receptors in long-term potentiation in the fimbria–CA3 pathway of the hippocampus
H. Katsuki, H. Saito and M. Satoh (Japan) 249
Cholinergic fiber loss occurs in the absence of synaptophysin depletion in Alzheimer's disease primary visual cortex
T.G. Beach and E.G. McGeer (Canada) 253
Evidence for increased calcium influx across the choroid plexus following brief ischemia of gerbil brain
J. Ikeda, G. Mies, T.S. Nowak Jr., F. Joó and I. Klatzo (USA) 257

*Acknowledgements to reviewers* 260

*Author Index* 275

*Erratum* 279